Available online at www.sciencedirect.com

# **ScienceDirect**

journal homepage: http://www.journals.elsevier.com/journal-of-palaeogeography/

Biopalaeogeography and palaeoecology

# A redescription of the ichnospecies *Koreanaornis anhuiensis* (Aves) from the Lower Cretaceous Qiuzhuang Formation at Mingguang city, Anhui Province, China



IOURNAL OF

PALAEOGEOGRAPHY

Li-Da Xing <sup>a,b,c</sup>, Yuan-Chao Hu<sup>d</sup>, Jian-Dong Huang <sup>d,\*</sup>, Qing He<sup>e</sup>, Martin G. Lockley <sup>f</sup>, Michael E. Burns <sup>g</sup>, Jun Fang <sup>b</sup>

- <sup>a</sup> State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Beijing 100083, China
- <sup>b</sup> School of Earth Sciences and Resources, China University of Geosciences, Beijing 100083, China
- <sup>c</sup> State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology. Chinese Academy of Sciences, Naniing 210008, Jiangsu Province, China
- <sup>d</sup> Anhui Geological Museum, Hefei 230031, Anhui Province, China
- <sup>e</sup> School of Resources and Environmental Engineering, Anhui University, Hefei 230039, Anhui Province, China
- <sup>f</sup> Dinosaur Trackers Research Group, University of Colorado Denver, PO Box 173364, Denver, CO 80217, USA
- <sup>g</sup> Department of Biology, Jacksonville State University, 700 Pelham Rd, Jacksonville, FL 36265, USA

**Abstract** The Cretaceous bird trackway originally labeled *Aquatilavipes anhuiensis*, in 1994, had previously been examined, photographed and replicated, but never described or illustrated in detail. However, it has been part of a widening discussion about the distribution of *Aquatilavipes* and *Koreanaornis* in China (and Korea). Here we illustrate and formally describe the holotype in detail and assign it to *Koreanaornis (Koreanaornis anhuiensis*) as informally proposed by previous authors. We also demonstrate that most authenticated reports of *Koreanaornis*, including the Anhui occurrence, are from the Lower Cretaceous, not from the Upper Cretaceous as previously reported.

**Keywords** Lower Cretaceous, Ichnospecies, *Koreanaornis, Aquatilavipes*, Qiuzhuang Formation, Anhui Province

© 2017 China University of Petroleum (Beijing). Production and hosting by Elsevier B.V. on behalf of China University of Petroleum (Beijing). This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Received 19 June 2017; accepted 18 July 2017; available online 6 December 2017



<sup>\*</sup> Corresponding author.

E-mail address: jiandongh2008@163.com (J.-D. Huang).

Peer review under responsibility of China University of Petroleum (Beijing).

https://doi.org/10.1016/j.jop.2017.10.003

<sup>2095-3836/© 2017</sup> China University of Petroleum (Beijing). Production and hosting by Elsevier B.V. on behalf of China University of Petroleum (Beijing). This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

### 1. Introduction

At present, many Early Cretaceous bird tracksite reports come from China (Lockley and Harris, 2010; Lockley et al., 2012a; Xing et al., 2011, 2015, 2016) and Korea (Huh et al., 2012; Lockley et al., 2012b). The number of new ichnogenera is steadily growing. In addition to Aquatilavipes (Zhen et al., 1994), the list now includes five recent additions: Shandongornipes (Li et al., 2005), Pullornipes (Lockley et al., 2006a), Wupus (Xing et al., 2007, 2015), Moguiornipes (Xing et al., 2011), Tatarornipes (Lockley et al., 2012c), and Uhangrichnus which, by formal revision, is considered a subjective senior synonym Dongyangornipes following Buckley et al. (2016), contra Azuma et al. (2013). Korea has also proved a rich source of Cretaceous bird tracksites including ichnospecies belonging to eight avian ichnogenera, including six first named from Korea (Kim et al., 2012, 2013; Lockley et al., 2012b; and references therein). Together China and Korea have yielded the vast majority of important Cretaceous bird tracks making East Asia a "paradise" for birds (Kim et al., 2012).

There were only two bird track records reported from China during 1990s, "Aquatilavipes sinensis" from Emei, Sichuan Province (Zhen et al., 1994), and "Aquatilavipes anhuiensis" from Mingguang, Anhui Province (Jin and Yan, 1994). Jin and Yan (1994) named "A. anhuiensis" based on a single trackway, which is the only specimen presently referred to this taxon. The specimen was mentioned by Matsukawa et al. (2006, p. 20) as "bird tracks" from the "Mingguang" (sic) site, and traced (UCM 569), photographed and replicated by one of us (MGL), in 2001, for the University of Colorado Museum (UCM) collections as UCM 214.45. Lockley et al. (2008, 2012a) suggested that "A. sinensis" had more affinity to Koreanaornis (Kim, 1969; Lockley and Harris, 2010; Lockley et al., 1992) than to "Aquatilavipes", and therefore reclassified it under the revised combination Koreanaornis sinensis. It is unlike Aquatilavipes in revealing a small hallux trace, and Lockley et al. (2013) also reclassified "A. anhuiensis" as Koreanaornis anhuiensis. We have since relocated the specimen allowing it to be described and illustrated here in detail based on independent examinations of the specimen more than a decade apart.

Jin and Yan (1994) did not mention geological setting or locality information in their brief paper describing the ichnotype of *K. anhuiensis*. Between 1999 and 2001, nothing was found during Sino-Japanese-American expeditions in Gupei Basin (also known as the Jiashan Basin) except for one potential site

( $32^{\circ}58'38.16''N$ ,  $117^{\circ}59'49.2''E$ ; Matsukawa *et al.*, 2006) from which the specimen may have originated. In the following years, the first author and researchers from Anhui Geological Museum also failed to make any progress in this area. However, further study of *K. sinensis* specimen and geological investigation of the site in 2009 and 2017 have provided new information that necessitates a revision of this specimen.

## 2. Geological setting

The Gupei Basin lies to the north of the Hefei Basin and is tectonically controlled by the extensional activities along the Tan–Lu Fault Zone like the Hefei Basin (Mercier *et al.*, 2007). The Yi–Shu Fault Belt extends from Shandong Province to Jiashan County in the north of Anhui Province with some Cretaceous graben basins developed along its extent. The Gupei Basin to the north of Jiashan County of Anhui Province is continuous with the Tancheng Basin in the south of Yi–Shu Fault Belt *via* Jiangsu Province (Zhu *et al.*, 2001).

Jin and Yan (1994) did not provide any geological background but only mentioned that "the fossil is found in Qiuzhuang Formation (formerly called Zhuxiang Formation) where plant fossils *Brachyphyllum*, *Otozamites*, *Suturovagina*, *Manica*, *Frenelopsis* with palynological assemblages dominated by *Schizaeoisporites* and other fossils such as bivalve *Sphaerium* had been discovered before, suggesting an early Late Cretaceous age" (Jin and Yan, 1994, p. 57).

However, Jin and Yan (1994) have confused some names of the formations. The Anhui Bureau of Geology and Mineral Resources (1987, 1988) considered that the Cretaceous sedimentary sequence of the Gupei Basin is divided into the Lower Cretaceous Xinzhuang Formation (formerly called Zhuxiang Formation), and Qiuzhuang Formation (formerly called Xiangdaopu Formation) and the Upper Cretaceous Zhangqiao Formation.

The Xinzhuang Formation is 1100 m in total thickness. The lower section of the Xinzhuang Formation mainly consists of gray purple conglomerate and gray yellow lithic sandstone interbedded with siltstone and mudstone. The upper section comprises yellow green medium-to fine-grained lithic sandstone, siltstone and mudstone interbedded with marlstone. This formation yields plant fossils (*Manica, Frenelopsis, Suturovagina,* and *Brachyphyllum*), bivalve (*Nakamuranaia chingshanensis*), conchostracan fossils (*Yanjiestheria* sp. and *Sinoestheria* sp.), and the ostracod *Cypridea* (*Cypridea*) priva Lubimova. Pollen of gymnosperms accounts for 81.3% of the palynological assemblage.



**Fig. 1** Geographical setting showing the location (footprint icon) of the Mingguang bird site in Anhui Province, and dinosaur sites in Shandong Province, China.

The largest proportion of pollen gymnosperms accounts is Pinaceae (25%), and secondly, *Classopollis* (18.6%). Most spores are attributable to *Lygodiaceae* (Anhui Bureau of Geology and Mineral Resources, 1987, 1988; Li and Jiang, 1997; Tu and Xiang, 1989).

The Qiuzhuang Formation crops out mostly in the Gupei and Hefei basins. The lower section is constituted of interbedded fine lithic sandstone, arkose sandstone and siltstone, intercalated with calcilutite and sandy mudstone. The lower part of the upper section is made up of interbedded gray purple thickbedded sandy conglomerate and fine-grained lithic arkose, and the upper part of the upper section is made up of interbedded dark purple silty mudstone and gray brown fine-grained lithic arkose. The total thickness is 2192 m, 900 m of which lies in the Hefei Basin and mainly crops out as brown and brownish gray silty mudstone, of which the upper part is interbedded with medium- to fine-grained sandstone that contains gypsum. The Qiuzhuang Formation is in conformable contact with the Xinzhuang Formation and preserves

similar plant fossils. In the palynological assemblage found in the Qiuzhuang Formation, spores account for 52.3%, and most of them are *Schizaeoisporites* (41.6%), and angiosperm pollen accounts for 3.6%— 6.1%. The ostracod *Cypridea* (*Pseudocypridina*) aversa and a few specimens of the bivalve *Sphaerium* can also be found in this formation (Anhui Bureau of Geology and Mineral Resources, 1987, 1988; Li and Jiang, 1997; Tu and Xiang, 1989).

Despite controversy over the geologic age of the Xinzhuang Formation (some authors proposed a Late Jurassic age (Zhang *et al.*, 2009)), the age of the Qiuzhuang Formation (Xiangdaopu Formation) is certainly the Early Cretaceous (Tu and Xiang, 1989; Zhang *et al.*, 2009; Zhu *et al.*, 2012). Zhu *et al.* (2012) suggested that the Xinzhuang Formation is Berriasian—Hauterivian in age, the Maotanchang Formation (volcanic rocks) is Barremian—Aptian, and the Qiuzhuang Formation is Albian based on crustal deformation kinematics and  ${}^{40}$ Ar/ ${}^{39}$ Ar dating. These formations are contemporaneous with the Dasheng, Qingshan

Table 1 Measurements (in cm) of the bird tracks from Mingguang site (Fig. 1), Anhui Province, China.													
Number	FL	FL′	FW	LI	LII	LIII	LIV	I—II	11—111	III—IV	PL	SL	L/W
AGB2882-1	2.32	_	4.06	_	2.14	2.30	2.20	_	61	77	_	_	0.6
AGB2882-2	2.76	3.49	3.12	0.76	1.55	2.68	2.10	101	51	61	5.33	-	0.9
AGB2882-3	<3.16	_	_	_	_	_	_	_	_	_	7.55	13.30	_
AGB2882-4	2.62	3.02	3.93	0.75	2.35	2.6	2.17	_	66	57	6.01	_	0.7
AGB2882-5	2.65	3.46	3.10	0.76	1.85	2.67	1.86	105	77	40	_	_	0.9
Mean	2.59	3.32	3.55	0.76	1.97	2.56	2.08	103	64	59	6.30	13.30	0.8

Abbreviations: FL = Footprint length; FL' = Footprint length with hallux; FW = Footprint width; LI, LII, LIII, LIV: Digit lengths; I–II, II–III, III–IV: Angle between digits I and II, digits II and IV; PL = Pace length; SL = Stride length; L/W = FL/FW.

(volcanic rocks) and Mengyin/Laiyang formations in the Yishu Graben and Jiaolai Basin, respectively.

## 3. Systematic ichnology

Koreanaornipodidae Lockley *et al.*, 2006b. *Koreanaornis* Kim, 1969.

# Type ichnospecies: *Koreanaornis hamanensis* Kim, 1969, Lockley *et al.*, 1992 emend.

**Diagnosis:** Small, tetradactyl bird tracks with small hallux impressions occasionally present. Digit traces typically separate (*i.e.*, not connected proximally). Claw traces variably present, slender, and obscure. Trackways exhibit inward (positive) rotation of the tracks. Tracks wider than long, with widths ranging from 2.5 cm to 4.4 cm. Divarication between digits II and IV averaging about 120° (Lockley *et al.*, 1992).

Koreanaornis anhuiensis n. comb (Jin and Yan, 1994). Lockley *et al.*, 2013.

Holotype: A complete natural cast of track from trackway with five steps cataloged as AGB2882-5 (GS VIII-4) from the Mingguang tracksite. The specimen is stored in the Anhui Paleontological Museum.

**Paratypes:** Track casts AGB2882-1-AGB2882-4 in the same trackway as the holotype stored in the Anhui Paleontological Museum.

**Type horizon and locality:** Qiuzhuang Formation, Lower Cretaceous. Mingguang tracksite, Anhui Province, China (Fig. 1).

**Emended diagnosis:** Small (footprint length ~33 mm, footprint width ~36 mm), sub-symmetrical, functionally tridactyl tracks by the following combination of characters: a consistently wide average digit II–IV divarication; consistently preserved, the midline of small hallux impression and digit III trace are aligned, and more than one third of the length of digit III trace; digit divarication I–II between 101° and 105° (Table 1; divarication II–IV between 112° and 138°).

Description (see Table 1 for reference): Small (3.32 cm average length; 3.55 cm average width) tetradactyl tracks, with small (0.75 cm-0.76 cm, n = 3) hallux impressions are consistently preserved, making the majority of the tracks appear functionally tridactyl. Digit I traces are relatively short compared to digit traces II-IV. Digit III is the longest digit, and digits II and IV are subequal in length. Digit II is broader than Digit IV. Digit traces II-IV connect proximally to a metatarsophalangeal pad. Digits II and III have two and three pads, respectively, and Digit IV pads are unidentifiable. Digit trace splay is high, with an average footprint length: width ratio of 0.8 (range of 0.6–0.9). The ratio of the length of the hallux impression to digit III is 1:3.5. Digit II–IV divarication (total divarication) is an average of  $123^{\circ}$  ( $112^{\circ}-138^{\circ}$  n = 4). The divarication of digits I-II is high (average 103°, range  $101^{\circ}-105^{\circ}$ , n = 2). The AGB2882 trackway is incomplete. Based on AGB2882-4 and AGB2882-5, the tracks exhibit an outward rotation.

Remarks: In all five tracks, only AGB2882-1 does not have hallux impression and is oriented in a different direction from the other tracks, indicating a different trackmaker. AGB2882-2-AGB2882-4 may, however, represent the same trackmaker, which turned the foot in an unusual way to make the impression of AGB2882-1 (Fig. 2). AGB2882-3 has distinct drag or sliding traces and double imprints of digits III and IV. Hallux impression of AGB2882-4 shows a lateral sliding trace. The characteristics in the well-preserved AGB2882-2 and AGB2882-5 are similar. Diagnostic characters proposed by Jin and Yan (1994) include similarities with characters suggested by Currie (1981) and by Zhen et al. (1994) with the exception of shorter digits and a  $128^{\circ}$ digit divarication between II and IV. Both digit length and digit divarication of AGB2882 fall into the range of Koreanaornipodidae.

**Discussion:** It is important to note that *Koreanaornis* is different from *Aquatilavipes*, in size and other morphological features (Buckley *et al.*, 2016). When *Aquatilavipes* was first reported from China by Zhen *et al.* (1994), these authors were unaware of the



**Fig. 2** Photograph (A), clean cast (B), 3D height map (warm colors are high, cooler colors are low) (C), contour map (D) and interpretative outline drawing (E) of *K. anhuiensis* AGB2882. A, C–E are in a same perspective. The rectangles in C and D are scale bar area from A.

original description of *Koreanaornis* from Korea (Kim, 1969). As a result, it was not until much later that it was recognized that such bird tracks, labeled as *Aquatilavipes* from China (Jin and Yan, 1994; Zhen *et al.*, 1994), were subjective junior synonyms of *Koreanaornis* (Buckley *et al.*, 2016; Lockley *et al.*, 2008, 2012a). However, it is still quite possible that *Aquatilavipes* may be found in China (Xing *et al.*, 2011).

Most characteristics of *K. anhuiensis* match those of Koreanaornipodidae (Kim, 1969; Lockley *et al.*,

2006b), such as small, wide, sub-symmetric, functionally tridactyl tracks with slender digit impressions and wide (90°–115°) divarication angles between digits II and IV. The published range of total divarication (digit divarication II–IV) for Koreanaornipodidae (Koreanaornis dodsoni, K. hamanensis, K. lii, Pullornipes aureus) is 32° (probably affected by muddy sediment)–170° (average 101°). With the exception of K. lii, the diagnostic orientation of digit I for Koreanaornipodidae is 180° from digit IV, which provides an



**Fig. 3** Morphologic comparison of the ichnospecies of Koreanaornipodidae. A and A' – *Koreanaornis hamanensis* (Lockley *et al.*, 2006b); B – *Koreanaornis lii* (Xing *et al.*, 2016); C and C' – *Koreanaornis sinensis* (Lockley *et al.*, 2008, 2012b); D – *Koreanaornis dodsoni* (Xing *et al.*, 2011); E – *Koreananornis anhuiensis*.

average digit I-II divarication of 79° (10°-148°). Both divarications of *K. anhuiensis* are in this range.

K. anhuiensis shares many features with K. hamanensis, K. dodsoni, K. lii and K. sinensis. However, significant differences exist. For example, K. hamanensis, K. dodsoni and K. lii have digit traces that do not join proximally on most tracks, but digit traces of K. anhuiensis connect proximally to a metatarsophalangeal pad. Digit trace slenderness in K. anhuiensis is more similar to K. hamanensis (Kim, 1969; Lockley *et al.*, 2006b, 2012b), K. sinensis (Lockley *et al.*, 2008; Zhen *et al.*, 1994) and P. aureus (Lockley *et al.*, 2006b) than to K. dodsoni (Xing *et al.*, 2011). Furthermore, K. dodsoni is larger in size, probably due to some degrees of preservational bias (Fig. 3).

The hallux impressions of *K. anhuiensis* tracks are only known from two specimens. Based on AGB2882-2 and AGB2882-5, the midline of the hallux impression almost overlaps that of digit III, different from other Koreanaornipodidae tracks. More specimens, however, are needed to determine the taxonomic significance of this feature.

#### 4. Conclusions

*Koreanaornis* is well known from tracksites in Korea. It was first discovered and named by Kim (1969) on the basis of abundant tracks from the Haman Formation (Aptian). Subsequently, the ichnotaxon was reported from the Jindong Formation (?Albian) of Goseong County (Lockley *et al.*, 1992) and other Korean localities (Kim, 2008). *Koreanaornis* was also recently described from the Early Cretaceous Tianjialou Formation (Barremian-Albian) Jishan tracksite in Shandong Province (Li *et al.*, 2005), on the basis of very small tracks (~2.5–3.0 cm wide and long). This occurrence indicates that *Koreanaornis* was also widely distributed in northern China in the late Early Cretaceous in addition to being abundant in Korea.

Based on the present knowledge, *Koreanaornis* is found in the late Early Cretaceous (Barremian—Albian) of China, as well as in deposits of broadly similar age in Korea (Lockley *et al.*, 2012b). This age range corresponds to the occurrence of *K. anhuiensis* from Anhui Province and supports the conclusion that it is inappropriate to consider these materials as Late Cretaceous (*contra* Jin and Yan, 1994).

#### Acknowledgements

The authors thank Yi-Ming Gong, Hong-Wei Kuang, Nasrollah Abbassi, Mike Benton and Bin Hu for critical comments and suggestions on this paper; Anthony Romilio, for preparing the 3D height maps; Shen-Na Wang, for the assistance in the field and measuring of the tracks. Martin G. Lockley had the opportunity to examine the *K. anhuiensis* specimen in 2001, thanks to the support of grant number 11791012 issued to Dr. Masaki Masukawa from Tokyo Gakugei University, Japan.

This research was supported by the National Natural Science Foundation of China (Grant No. 41772008), the State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences (Grant No. 173127), and the Special Scientific Research Fund of the Non-Profit Sector Project, Ministry of Land and Resources of China (Grant No. 201511054).

#### References

- Anhui Bureau of Geology and Mineral Resources, 1987. *Regional Geology of Anhui Province*. Geological Publishing House, Beijing, 470 pp. (in Chinese).
- Anhui Bureau of Geology and Mineral Resources, 1988. Stratigraphic Memoirs of Anhui (Cretaceous Volume and Jurassic Volume). Anhui Science and Technology Press, Hefei, pp. 66–126 (in Chinese).
- Azuma, Y., Lü, J.C., Jin, X.S., Noda, Y., Shibata, M., Chen, R.J., Zheng, W.J., 2013. A bird footprint assemblage of early late Cretaceous age, Dongyang City, Zhejiang Province, China. *Cretaceous Research*, 40, 3–9.
- Buckley, L.G., McCrea, R.T., Lockley, M.G., 2016. Analysing and resolving Cretaceous avian ichnotaxonomy using

multivariate statistics: Approaches and results. In: Falkingham, P.L., Marty, D., Richter, A. (Eds.), *Dinosaur Tracks: The Next Steps.* Indiana University Press, pp. 258–308.

- Currie, P.J., 1981. Bird footprints from the Gething formation (Aptian Lower Cretaceous) of northeastern British Columbia, Canada. *Journal of Vertebrate Paleontology*, 1(3-4), 257–264.
- Huh, M., Lockley, M.G., Kim, K.-S., Kim, J.-Y., Gwak, S.-G., 2012. First report of *Aquatilavipes* from Korea: New finds from Cretaceous strata in the Yeosu Islands Archipelago. *Ichnos*, 19, 43–49.
- Jin, Y., Yan, H., 1994. Bird footprints discovered in the Cretaceous red bed of the Gupei Basin, Anhui. *Geology* of Anhui, 4(3), 57–61 (in Chinese with English abstract).
- Kim, B.K., 1969. A study of several sole marks in the Haman Formation. *Journal of the Geological Society of Korea*, 5, 243–258.
- Kim, J.-Y., Kim, M.K., Seop Ho, M., Zin Lee, C., 2013. A new semi-palmate bird track, *Gyeongsangornipes lockleyi* ichnogen. et ichnosp. nov., and Koreanornis from the Early Cretaceous Jindong Formation of Gosong County, southern coast of Korea. *Ichnos*, 20, 72–80.
- Kim, J.-Y., Seo, S.J., Kim, K.S., Lockley, M.G., Kim, S.H., Baek, K.S., 2012. A paradise of Mesozoic birds: The world's richest and most diverse Cretaceous bird track assemblage from the Early Cretaceous Haman Formation of the Gajin tracksite, Jinju, Korea. *Ichnos*, 19, 28–42.
- Kim, S.-H., 2008. Stratigraphic Diversity of Bird and Diminutive Dinosaur Tracks from the Cretaceous Haman Formation, South Gyeongsang Province of Korea: Implication for Ichnostratigraphy. Ph.D. Dissertation. Graduate School of Korea National University of Education, 141 pp.
- Li, R.H., Lockley, M.G., Liu, M.W., 2005. A new ichnotaxon of fossil bird track from the Early Cretaceous Tianjialou Formation (Barremian—Albian), Shandong Province, China. *Chinese Science Bulletin*, 50, 1149—1154.
- Li, Y.F., Jiang, L.F., 1997. Stratigraphy (Lithostratic) of Anhui Province (in Chinese). China University of Geosciences Press, Wuhan, pp. 1–271.
- Lockley, M.G., Harris, J.D., 2010. On the trail of early birds: A review of the fossil footprint record of avian morphological and behavioural evolution. In: Ulrich, P.K., Willett, J.H. (Eds.), *Trends in Ornithology Research*. Nova Science Publishers, Inc., Hauppauge, pp. 1–63.
- Lockley, M.G., Houck, K., Yang, S.-Y., Matsukawa, M., Lim, S.-K., 2006b. Dinosaur dominated footprint assemblages from the Cretaceous Jindong Formation, Hallyo Haesang National Park area, Goseong County, South Korea: Evidence and implications. *Cretaceous Research*, 27, 70–101.
- Lockley, M.G., Kim, J.Y., Kim, K.S., Kim, S.H., Matsukawa, M., Li, R., Li, J., Yang, S.-Y., 2008. *Minisauripus* — The track of a diminutive dinosaur from the Cretaceous of China and South Korea: Implications for stratigraphic correlation and theropod foot morphodynamics. *Cretaceous Research*, 29, 115–130.
- Lockley, M.G., Li, J.J., Li, R.H., Matsukawa, M., Harris, J.D., Xing, L.D., 2013. A review of the tetrapod track record in

China, with special reference to type ichnospecies: Implications for ichnotaxonomy and paleobiology. Acta Geologica Sinica (English Edition), 87, 1-20.

- Lockley, M.G., Li, J.J., Matsukawa, M., Li, R., 2012c. A new avian ichnotaxon from the Cretaceous of Nei Mongol, China. *Cretaceous Research*, 34, 84–93.
- Lockley, M.G., Lim, J.D., Kim, J.Y., Kim, K.S., Huh, M., Hwang, K.-G., 2012b. Tracking Korea's early birds: A review of Cretaceous avian ichnology and its implications for evolution and behavior. *Ichnos*, 19, 17–27.
- Lockley, M.G., Lucas, S.G., Matsukawa, M., Harris, J.D., 2012a. Cretaceous tetrapod footprint biostratigraphy, biochronology, and ichnofacies. *Journal of Stratigraphy*, 36, 503–551.
- Lockley, M.G., Matsukawa, M., Ohira, H., Li, J.J., Wright, J., White, D., Chen, P.J., 2006a. Bird tracks from Liaoning Province, China: New insights into avian evolution during the Jurassic-Cretaceous transition. *Cretaceous Research*, 27, 33–43.
- Lockley, M.G., Yang, S.Y., Matsukawa, M., Fleming, F., Lim, S.K., 1992. The track record of Mesozoic birds: Evidence and implications. *Philosophical Transactions of the Royal Society of London B*, 336, 113–134.
- Matsukawa, M., Lockley, M., Li, J., 2006. Cretaceous terrestrial biotas of East Asia, with special reference to dinosaur-dominated ichnofaunas: Towards a synthesis. *Cretaceous Research*, 27, 3–21.
- Mercier, J.L., Hou, M.J., Vergely, P., Wang, Y.M., 2007. Structural and stratigraphical constraints on the kinematics history of the southern Tan—Lu Fault Zone during the Mesozoic Anhui Province, China. *Tectonophysics*, 439, 33–66.
- Tu, Y.J., Xiang, J.X., 1989. Tripartite division of the Cretaceous and the Cretaceous-Jurassic boundary in Anhui. *Geological Bulletin of China*, 1, 28–35 (in Chinese with English abstract).
- Xing, L.D., Buckley, L.G., Lockley, M.G., Zhang, J.P., Marty, D., Wang, Y., Li, J.J., McCrea, R.T., Peng, C., 2016. A new bird track, *Koreanaornis lii* ichnosp. Nov., from the lower Cretaceous Hekou Group in the Lanzhou-Minhe Basin, Gansu, Northwest China, and implications for early Cretaceous avian diversity. *Cretaceous Research*, 66, 141–154.
- Xing, L.D., Buckley, L.G., McCrea, R.T., Lockley, M.G., Zhang, J.P., Piñuela, L., Klein, H., Wang, F.P., 2015. Reanalysis of Wupus agilis (Early Cretaceous) of Chongqing, China as a large avian trace: Differentiating between large bird and small non-avian theropod tracks. *PLoS One*, 10(5), e0124039.
- Xing, L.D., Harris, J.D., Jia, C.K., Luo, Z.J., Wang, S.N., An, J.F., 2011. Early Cretaceous bird-dominated and dinosaur footprint assemblages from the Northwestern Margin of the Junggar Basin, Xinjiang, China. *Palaeoworld*, 20, 308–321.
- Xing, L.D., Wang, F.P., Pan, S.G., Chen, W., 2007. The discovery of dinosaur footprints from the Middle Cretaceous Jiaguan formation of Qijiang County, Chongqing City. *Acta Geologica Sinica (Chinese Edition)*, 81(11), 1591–1602 (in Chinese with English abstract).
- Zhang, J.D., Hao, T.Y., Fan, D.H., Liu, D.L., Zhao, Y.H., 2009. Identification of geological time of Mesozoic strata in Hefei Basin and its significance. *Acta Geologica Sinica*

(Chinese Edition), 83(5), 599–608 (in Chinese with English abstract).

- Zhen, S.N., Li, J.J., Chen, W., Zhu, S., 1994. Dinosaur and bird footprints from the lower Cretaceous of Emei County, Sichuan. *Memoirs of the Beijing Natural History*, 54, 105–120 (in Chinese with English abstract).
- Zhu, G., Jiang, D.Z., Zhang, B.L., Chen, Y., 2012. Destruction of the eastern North China Craton in a backarc

setting: Evidence from crustal deformation kinematics. *Gondwana Research*, 22, 86–103.

Zhu, G., Wang, D.X., Liu, G.S., Song, C.Z., Xu, J.W., Niu, M.L., 2001. Extensional activities along the Tan–Lu fault zone and its geodynamic setting. *Chinese Journal of Geology*, 36(3), 269–278 (in Chinese with English abstract).